


AD-A193 520


ANALYSIS OF RADIATION EXPOSURE - SERVICE PERSONNEL ON
RONGERIK ATTLL: OPE (U) SCIENCE APPLICATIONS
INTERNATIONAL CORP MCLEARN VA J GOETZ ET AL 09 JUL 87
SATC-86/1608 DNA-TR-86-120 DNA001-85-C-0101 F/G 6/7

1/1

UNCLASSIFIED

NL





DNA-TR-86-120

AD-A193 520

**ANALYSIS OF RADIATION EXPOSURE —
SERVICE PERSONNEL ON RONGERIK ATOLL
Operation Castle - Shot Bravo**

J. Goetz, et al.
Science Applications International Corp.
P.O. Box 1303
McLean, VA 22102-1303

9 July 1987

Technical Report

CONTRACT No. DNA 001-85-C-0101

Approved for public release;
distribution is unlimited.

THIS WORK WAS SPONSORED BY THE DEFENSE NUCLEAR AGENCY
UNDER RDT&E RMC CODE B350084466 RM RH 00012 25904D.

DTIC
ELECTE
APR 19 1988
S H D

Prepared for
Director
DEFENSE NUCLEAR AGENCY
Washington, DC 20305-1000

88

Destroy this report when it is no longer needed. Do not return to sender.

PLEASE NOTIFY THE DEFENSE NUCLEAR AGENCY
ATTN: TITL, WASHINGTON, DC 20305 1000, IF YOUR
ADDRESS IS INCORRECT, IF YOU WISH IT DELETED
FROM THE DISTRIBUTION LIST, OR IF THE ADDRESSEE
IS NO LONGER EMPLOYED BY YOUR ORGANIZATION.



DISTRIBUTION LIST UPDATE

This mailer is provided to enable DNA to maintain current distribution lists for reports. We would appreciate your providing the requested information.

☐ Add the individual listed to your distribution list.

☐ Delete the cited organization/individual.

☐ Change of address.

NAME: _____

ORGANIZATION: _____

OLD ADDRESS

CURRENT ADDRESS

_____	_____
_____	_____
_____	_____

TELEPHONE NUMBER: (____) _____

_____	_____
_____	_____
_____	_____

SUBJECT AREA(S) OF INTEREST: _____

DNA OR OTHER GOVERNMENT CONTRACT NUMBER: _____

CERTIFICATION OF NEED-TO-KNOW BY GOVERNMENT SPONSOR (if other than DNA): _____

SPONSORING ORGANIZATION: _____

CONTRACTING OFFICER OR REPRESENTATIVE: _____

SIGNATURE: _____

CUT HERE AND RETURN



Director
Defense Nuclear Agency
ATTN: TITL
Washington, DC 20305-1000

Director
Defense Nuclear Agency
ATTN: TITL
Washington, DC 20305-1000

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No 0704 0188 Exp Date Jun 30, 1986	
1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY N/A since Unclassified			3. DISTRIBUTION AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A since Unclassified					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) SATC-86/1006			5. MONITORING ORGANIZATION REPORT NUMBER(S) DNA-TR-86-120		
6a. NAME OF PERFORMING ORGANIZATION Science Applications International Corp.		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Director Defense Nuclear Agency		
6c. ADDRESS (City, State, and ZIP Code) P.O. BOX 1303 McLean, Virginia 22102-1303			7b. ADDRESS (City, State, and ZIP Code) Washington, DC 20305-1000		
8a. NAME OF FUNDING SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable) STBE/Auton	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DNA 001-85-C-0101		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO 62715H	PROJECT NO RM	TASK NO RH WORK UNIT ACCESSION NO DH009019
11. TITLE (Include Security Classification) ANALYSIS OF RADIATION EXPOSURE - SERVICE PERSONNEL ON RONGERIK ATOLL Operation Castle- Shot Bravo					
12. PERSONAL AUTHOR(S) Metz, J.; Klemm, J.; Phillips, J.; Thomas, C.					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 850312 TO 870612		14. DATE OF REPORT (Year, Month, Day) 870709	
15. PAGE COUNT 48					
16. PRELIMINARY NOTATION This work was sponsored by the Defense Nuclear Agency under RDT&E RMC Code B350084466 RM PH 00012 25904D.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB GROUP	Operation CASTLE, Shot Bravo Rongerik Atoll		
07	07		Radiation Exposure Assessment		
			Nuclear Test Personnel Review (NTPR)		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) External and internal doses are reconstructed for the 28 American servicemen stationed on Rongerik Atoll, Marshall Islands, who were exposed to fallout on 1-2 March 1954 from Shot Bravo of Operation CASTLE. External doses are determined from limited radiation survey and film badge information. Internal dose commitments are derived from urinalysis data. The magnitude of the calculated activity intake suggests the principal pathways. Reconstructed film badge doses are approximately 40 rem, with adjustments from individual activity monitoring, as available. Internal dose commitments to the thyroid and large intestine (mainly all first-year dose) provide the only significant increments to the external dose. Total doses are approximately 230 rem to the thyroid, 115 rem to the lower large intestine, 30 rem to the upper large intestine, and about 40 to 50 rem to all other organs.					
20a. TYPE AND EXTENT OF ABSTRACT <input type="checkbox"/> FULL ABSTRACT <input checked="" type="checkbox"/> SUMMARY <input type="checkbox"/> EXTENDED SUMMARY			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. AUTHOR (Last Name, First Name, Middle Initial) Young, J. L.			22b. TELEPHONE (Include Area Code) (202) 325-7042	22c. OFFICE SYMBOL DNA/CSTI	


DD FORM 1473, 1-79

B3 APP edition may be used until exhausted
All other editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE



UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

TABLE OF CONTENTS

Section		Page
	LIST OF TABLES	iv
1	INTRODUCTION AND SUMMARY	1
2	BACKGROUND AND AVAILABLE DATA	2
	2.1 BACKGROUND	2
	2.2 AVAILABLE DATA AND ASSUMPTIONS	3
3	TECHNICAL APPROACH -- EXTERNAL DOSE	6
	3.1 DOSE CALCULATIONS	6
	3.2 AVAILABLE FILM BADGE DATA	12
	3.3 COMPARISON OF ESTIMATED FILM BADGE DOSES WITH FILM BADGE DATA	14
4	TECHNICAL APPROACH -- INTERNAL DOSE	16
	4.1 RELATIVE ACTIVITY	17
	4.2 INVENTORY NORMALIZATION	17
	4.3 PATHWAYS AND DOSE CONVERSION FACTORS	19
	4.4 ORGAN-SPECIFIC DOSE COMMITMENTS	22
5	RESULTS AND COMPARATIVE ANALYSIS	23
	5.1 RESULTS	23
	5.2 COMPARISON WITH PREVIOUS DOSE DETERMINATIONS	26
	5.3 TOTAL DOSE	30
6	LIST OF REFERENCES	31
Appendix	SAMPLE CASE INTERVIEW	35

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

LIST OF TABLES

Table		Page
1	Preliminary dose estimates for personnel on Rongerik Atoll, 1-2 March 1954.	11
2	Film badge dosimetry results, Rongerik Atoll, 1-2 March 1954.	14
3	Comparison of calculated film badge doses with dosimetry.	15
4	Reconstructed gamma radiation doses for military personnel stationed on Rongerik Atoll, 1 and 2 March 1954.	24
5	Fifty-year internal dose commitments for military personnel stationed on Rongerik Atoll, 1 and 2 March 1954.	25
6	Calculated film badge dose for one airman (CN 406).	56

SECTION I

INTRODUCTION AND SUMMARY

During the period 1-2 March 1954, twenty-eight Army and Air Force personnel stationed at Rongerik Atoll in the Marshall Islands were exposed to radioactive fallout from Shot Bravo of Operation CASTLE. The purpose of this report is to determine the whole body gamma radiation dose received by these personnel prior to their evacuation from the atoll, plus their organ dose commitments from internal emitters.

The external gamma radiation dose (i.e., dose equivalent) is reconstructed as follows. The radiological environment resulting from Shot Bravo fallout is determined from time-of-arrival data on Rongerik, modeled deposition, measured decay rates, and a later reading on Rongerik. Personnel activity scenarios are considered in conjunction with radiation shielding factors to arrive at film badge doses on an individual basis. Although film badge readings are available, they relate inadequately to personnel dose. Readings that reflect well-specified exposures indicate good agreement with corresponding calculated doses. A small, but systematic, difference suggests an adjustment to the calculated personnel doses, which are in the vicinity of 40 rem.

Organ-specific 50-year dose commitments (i.e., committed dose equivalent) resulting from the large-particle inhalation and ingestion of Bravo fallout radionuclides are calculated. Intake of iodine-131 is determined from the body metabolism of iodine that led to a measured activity of I-131 in a pooled urine sample taken 17 days after Bravo. The radionuclide inventory for the time of intake is normalized to the level of I-131 activity, thereby indicating the full intake of radioactivity. Large-particle inhalation dose conversion factors for each radionuclide are developed for calculation of internal dose commitments to each organ. The calculated thyroid dose is 190 rem, and dose to the intestines is as great as 76 rem. Other organ doses do not add significantly to the external gamma dose.

SECTION 2

BACKGROUND AND AVAILABLE DATA

2.1 BACKGROUND.

Shot Bravo was detonated on Bikini Atoll at 0645 hours (local time) on 1 March 1954. The total yield of the thermonuclear explosion was approximately 15-megatons TNT equivalent. The nuclear device was mounted on a barge in shallow water, about seven feet above the surface of a coral reef. The radioactive fallout from Shot Bravo contaminated an area extending from about 20 statute miles upwind to over 330 miles downwind and varying in width to over 60 miles. Included within the area of major contamination were Bikini Atoll and three downwind atolls: Ailinginae Atoll, Rongelap Atoll, and Rongerik Atoll (References 1, 2). Among these atolls, Marshallese were present on Ailinginae and Rongelap, and American servicemen only on Rongerik.

Twenty-eight servicemen (25 Air Force personnel and 3 Army personnel) operated a weather station on Rongerik Atoll, about 150 miles east of Bikini. At approximately 1407 hours (almost 7½ hours after the Bravo detonation) the deposition of fallout on Rongerik was detected by a low-level gamma radiation monitoring instrument and subsequently observed visually.

At 1500 hours, a message was sent to Joint Task Force (JTF) Headquarters on Enewetak Atoll (Reference 3) notifying them of the fallout on Rongerik. Approximately one-half hour later, a reply came back from Enewetak instructing the military personnel on Rongerik to put on long-sleeved shirts, trousers, hats, and GI shoes, and to remain inside as much as possible, consistent with their normal work routine. At about 2330 hours, another message was received directing all personnel to cease operations and move inside permanent buildings.

At 1245 hours on 2 March 1954, eight men, the first eight on the alphabetical detachment roster, were evacuated from Rongerik to Kwajalein Atoll by airplane; the remaining twenty men were evacuated by airplane at approximately 1800 hours (Reference 3).

Upon arrival at Kwajalein Atoll, the personnel evacuated from Rongerik were monitored for personal contamination and decontaminated. The decontamination program, consisting of repeated showers and radiation monitoring, continued from 2 March through 6 March. While at Kwajalein, the film badges that had originally been issued to the Rongerik detachment were collected and processed. In addition, urine samples, both pooled and individual, were collected and sent to laboratories in the United States for analysis. Late in April, the servicemen from Rongerik were transferred from Kwajalein to Tripler Army Hospital (Honolulu, Hawaii) for further observation and subsequent return to duty (References 3, 4).

2.2 AVAILABLE DATA AND ASSUMPTIONS.

Just prior to the first evacuation on 2 March, a gamma intensity reading was obtained with an uncalibrated AN/PDR-39 radiation survey meter. Because the operating condition of the instrument was not known at the time of its use, this reading (2000 mR/hr @ H+28.5 hours--Reference 5) is used only to indicate the general magnitude of the fallout intensity at the time of the first evacuation. The first radiological survey with calibrated instruments on the atoll was conducted on 10 March 1954, nine days after the shot. At this time the rad-safe survey team encountered average outdoor radiation intensity readings of 280 mR/hr on the island where the military personnel had been stationed. Other survey readings on 10 March were as discussed in Section 3.1. Additional surveys were made in subsequent weeks, but the readings were not as comprehensive and had less internal consistency. This likely arose from differential weathering and decay. Therefore, these later readings are not used in this analysis.

The only other radiation intensity data available from Rongerik were obtained on 1 March when a low-level gamma background monitoring instrument at the weather station began to register at 1407 hours and then went off scale (100 mR/hr) at 1437 (H+7.2 hr). The data from this instrument establish the time of arrival of the fallout.

After Shot Bravo, analyses of fallout samples were made to determine the decay of gamma intensity with time. Unfortunately, the only data adequate to characterize

the fallout were from Bikini Atoll and may not be completely representative of the fallout decay on Rongerik Atoll. Radiation intensity readings obtained from the Bikini lagoon (How Island) indicated decay rates that varied considerably from the traditional $t^{-1.2}$ rule. Average values for the decay exponent, obtained with several gamma ionization time-intensity meters on Bikini (Reference 6), are as follows:

$$\begin{aligned} 3 \leq t < 10 \text{ hrs} \quad k &= -1.19 \\ 10 \leq t < 48 \text{ hrs} \quad k &= -0.815 \\ 48 \leq t < 480 \text{ hrs} \quad k &= -1.50 \end{aligned}$$

Figure 1 depicts this fallout decay in comparison to $t^{-1.2}$ decay. A varying decay of this type is consistent with the presence of Np-239 ($t_{1/2}=56$ hr) and U-240 ($t_{1/2}=14$ hr), significant neutron-activation products of the U-238 in a thermonuclear device.

Because fallout decay rate data on Rongerik Atoll are not available, dose calculations for the military personnel on Rongerik are made using the measured decay rates on Bikini and the measured radiation intensity on Rongerik Atoll (280 mR/hr at D+9 days). In addition, the radiation environment during fallout deposition must be more explicitly defined. From the gamma background monitoring instrument on Rongerik, the time of arrival of measurable fallout from Bravo has been determined to be 1407 hours, or 7.4 hours after the shot. The radiation time-intensity recordings on Bikini indicated that, once fallout began, a period of 1-2 hours elapsed while fallout was occurring before the radiation levels reached a maximum and began to decay, even though fallout was still continuing (Reference 7). Other data were obtained aboard three Task Force 7.3 ships that were positioned at various distances downwind from the Bravo GZ. On one of these ships--the GYPSY--fallout commenced approximately 7 hours after detonation, about the time of arrival on Rongerik. The GYPSY data indicated that at least 4 hours of significant fallout occurred. The exact duration cannot be determined because ship washdown procedures were initiated while the intensity was still increasing. The data indicate, however, that at locations downwind from Bikini, the time to maximum intensity was somewhat longer than it was on Bikini. Based on these data, a five-hour period of significant fallout deposition is used to characterize the radiation environment on Rongerik. It is assumed that 90 percent of the fallout, by mass, was deposited during this interval.




Figure 1. Fallout intensity versus time after burst (Shot Bravo) normalized to intensity reading on Rongerik.

SECTION 3

TECHNICAL APPROACH--EXTERNAL DOSE

3.1 DOSE CALCULATIONS.

In order to calculate the radiation dose (i.e., dose equivalent) to personnel on Rongerik from the Shot Bravo fallout, the free-field radiation environment must be specified mathematically. Further, the calculated doses are dependent on any protection or shielding that was provided by buildings in which they stayed until they were evacuated.

(1) Free-Field Radiation on Rongerik

Based on the available data, the free-field radiation environment on Rongerik is estimated as follows:

For times greater than 48 hours after burst, the radiation decay followed a $t^{-1.5}$ decay rate. From the radiation intensity of 0.28 R/hr measured on Rongerik on 10 March (about 216 hours after burst), the intensity at 48 hours after burst (I_{48}) is:

$$I_{48} = I_{216} (216/48)^{1.5} = 2.67 \text{ R/hr} \quad (1)$$

Similarly, from I_{48} and the radiation decay according to $t^{-0.815}$ between 10 and 48 hours after detonation, the intensity after cessation of fallout through the time of evacuation is:

$$I_t = I_{48} (48/t)^{0.815} = 2.67 (48/t)^{0.815} = 62.7 t^{-0.815} \quad (2)$$

During fallout deposition, this decay applies to landed fallout particles (the change in decay exponent at 10 hours is too early in the deposition to noticeably affect the calculations). Thus, it is necessary to determine the fraction of time-normalized activity (or mass, for constant specific activity) deposited with time. This deposition is assumed to occur as a Gaussian (normal) distribution in time, reflecting a similar

spatial (horizontal) distribution in the nuclear cloud. A correction is applied for lateral diffusion of the cloud. For an increase in cloud area in direct proportion to time, the concentration of fallout particles is inversely proportional to time. The rate of deposition with time (T) is of the form:

$$cT^{-1}e^{-a(T-b)^2}$$

Parameters a and b are chosen to match the meter reading of 100 mR/hr at 7.9 hours and for the central 90 percent of the deposition to occur in 5 hours. The normalization to total deposition is provided by c. Fallout accumulated through time t is represented by the time integral of the above expression. The choice of the lower limit of integration is not crucial; the observed time of fallout onset is satisfactory. The complete expression for intensity with time is then:

$$I_t = 62.7t^{-0.815} \cdot 3.09 \int_{7.4}^t T^{-1} e^{-0.23(T-11.6)^2} dT \quad (3)$$

Figure 2 depicts this relationship, which serves as the basis for the dose calculations. The peak intensity of 7 R/hr occurs at the end of the estimated interval of significant fallout (9 to 14 hours).

(2) Personnel Activities on Rongerik Atoll, 1-2 March 1954

After being evacuated to Kwajalein, each person who was on Rongerik was interviewed to determine his specific activities during the period 1 March until evacuation on 2 March. Other pertinent data concerning their exposure conditions were also obtained such as clothing worn, food and drink consumed, and their recollection of what the fallout resembled. A synopsis of all of these interviews (Reference 3) is used in reconstructing the radiation exposure estimates in this memorandum. Seventeen of the interviews were in sufficient detail that the individuals' activities during the exposure period could be fairly well established on an hour-to-hour basis. For intervals without specified activities, reasonable assumptions are made that high-side the calculated radiation doses. The Appendix contains an example of the information available and the procedure employed in order to reconstruct the dose for one individual.




Figure 2. Rongerik fallout intensity versus time after burst (Shot Bravo).

(3) Protection Factors on Rongerik Atoll

During the initial radiological survey of Rongerik on 10 March 1954 (Reference 8), specific locations were surveyed with an AN/PDR-39 survey meters. In addition to the average reading of 280 mR/hr found at various locations on the island, the survey team also took intensity readings both inside and outside of the various structures utilized by the military personnel during their exposure period. The ratio of the intensity outside to the radiation intensity inside is defined as the protection factor and gives a measure of the radiation protection or shielding provided by the structure. From the 10 March survey data, each type of structure and its measured protection factor are as follows:

<u>Structure</u>	<u>Protection Factor</u>
Mess Hall	2.0
Tent	1.5
Latrine	1.6
Sleeping quarters	2.9
Dispensary	2.0
R-Section Bldg	2.5

(4) Film Badge Doses

With the free-field radiation intensity (I_t) as defined previously, film badge doses for the military personnel on Rongerik can be calculated for each time interval (t_1 to t_2) to which a protection factor (PF_i) applies, as follows:

$$D_{fb} = (0.7)(1/PF_i) \int_{t_1}^{t_2} I_t dt \quad (4)$$

where: D_{fb} = Dose as would be recorded by a film badge worn on the chest.
(0.7) = Factor to convert integrated free-field radiation intensity to film badge dose. This factor takes into account the body shielding of the badge as worn.

The maximum possible dose to a film badge worn continuously through the time of final evacuation is obtained by using $PF=1$ throughout, i.e., no radiation protection available on the island. The upper limit of integration in this case would be 35 1/4 hours after burst, which corresponds to final evacuation at 1800 hours, 2 March. A dose of 87 rem results. If a film badge was not worn, but instead left out in the open, the (0.7) factor would not enter into the calculations. As an example, one film badge was left hanging from a tent post (PF assumed to be 1) until H+28.75 hours (1130 hours, 2 March) at which time it was placed in the mess hall ($PF=2$) until the second group evacuated. The calculated dose is 112 rem.

All of the film badge dose calculations are dependent on the accuracy of integrated intensities derived from Figure 2. Uncertainty, especially concerning the duration of fallout deposition, is evidenced in a comparison with limited film badge dosimetry data (Section 3.3). Thus, film badge dose calculations in this section represent initial estimates.

For the seventeen individuals who provided sufficient information in their post-evacuation interviews to reconstruct their activities, the preliminary film badge dose estimates are given in Table 1. Because the individual activities used to derive these film badge doses were varied enough to encompass most reasonable activities, the calculations can be applied with confidence to all individuals, even those with insufficient information on their specific activities.

As would be expected, the second group of evacuees received a greater film badge dose than did the group evacuated approximately 5 hours earlier. Further, among the second group of evacuees, it appears that the Army troops who were manning the Project 6.6 station on Rongerik received a slightly larger dose than did the Air Force troops. This is not unexpected since the Army area on the island did not have any "permanent" structures, and the troops slept in a tent until approximately midnight, 1 March.

Table 1. Preliminary dose estimates for personnel on Rongerik Atoll, 1-2 March 1954.

Clinical No. (CN) <u>Assigned**</u>	Calculated Film Badge Dose (rem)	
	<u>Air Force</u>	<u>Army</u>
First Group to Evacuate		
401	*	
402	*	
403	*	
404	39	
405	*	
406	35	
407	35	
408	*	
Second Group to Evacuate		
409	46	
410	*	
411	46	
412	38	
413	*	
414		51
415	43	
416	43	
417	44	
418	41	
419	*	
420	42	
421	*	
422	40	
423		48
424	*	
425		45
426	38	
427	44	
428	*	

*Film badge dose is assumed comparable to that dose received by other personnel in the evacuation group.

**For this analysis, clinical numbers are used instead of actual names.

3.2 AVAILABLE FILM BADGE DATA.

Film badge data pertaining to the radiation exposure on Rongerik are reported in several references; however, the readings require interpretation. Because the allotted badges were not worn continuously by each individual, the data do not necessarily reflect the doses accrued by the servicemen. Moreover, there are inconsistencies among the references regarding the assignment of badges and in some of the reported readings.

Either twelve (Reference 9) or fourteen (Reference 3) film badges were received at the Rongerik weather station on 19 February 1954. One of these (#314) was distributed to the Army group. It was positioned seven feet above the floor of a tent in the Army area and was not worn. Two badges (#310/311) were assigned to the radio-weather station and were worn by a member of each shift (CN 408/CN 416) until approximately 1130 hours, 2 March. One badge (#312) was to be worn in the housing area by one of the Air Force personnel (CN 411) and another badge (#313), was worn by the NCOIC of the camp (CN 419). One other badge (#315) was placed on the side of a building in the housing area. This one was located shoulder high and moved occasionally each day to avoid direct sunlight. The remaining six badges (#309, #316-#320) were placed in a refrigerator in the mess hall. According to Reference 3, there were eight film badges placed in the refrigerator, but the reference does not cite badge numbers as does Reference 9.

Of the six badges that were issued, five (all but #313) were collected at 1130 hours, 2 March, and placed in a manila envelope and set on a table in the mess hall. The film badges that had been in the refrigerator remained there until 1700, at which time they too were placed in the manila envelope in preparation for the final evacuation. CN 419 continued to wear badge #313 until he arrived at Kwajalein at 1900, 2 March 1954.

The first written report of the film badge doses accrued on Rongerik came in Dispatch No. 240 431A from CTG 7.1, Eniwetok and COMNAVSTA, Kwajalein, received 25 April 1954, which stated (documents of the period reported doses in r (i.e., R), not the modern rem):

Film Badge Readings are as follows:

CN 401	40 r	CN 416	40 r
CN 403	40 r	CN 417	40 r
CN 404	40 r	CN 419	52 r
CN 408	40 r	Unknown	40 r
CN 410	44 r	CN 414	98 r
CN 411	40 r	CN 423	98 r
CN 415	40 r	CN 425	98 r

Film badges presently in custody of TU 7, TG 7.1 until completion of operation, at which time they will be forwarded to AFSWP.

This dispatch reportedly (Reference 3) contains the distribution list for the badges as originally planned; however, because only six of the badges were actually distributed, and only four were actually worn, many of the 40 R readings attributed to personnel may have actually been from the film badges that were placed in the refrigerator. One individual with whom a badge reading is associated was not stationed on Rongerik during the period 1-2 March 1954.

Reference 10 states that: "Several badges worn both outdoors and inside buildings on the island read 50-65 r, and one badge which remained outdoors over the 28.5 hr period read 98 r. Another group kept indoors inside a refrigerator read 38 r." The inference is that the troops received a film badge dose ranging from 50-65 R, while the badges in the refrigerator and the one that remained outside provide a lower and upper bound dose, respectively.

Reference 5, which cites a Hq JTF-7 Memo for Record: "Rad Safe Narrative Sequence of Events", BRAVO Shot, Operation CASTLE, as a reference, states that "film badge readings were obtained covering a range of values which varied with exposure conditions. Several badges were worn both outdoors and indoors (and covered the range 44-52 r). One badge which remained outdoors over the 28.5 hour exposure reached the upper limit of 98 r. Several other badges kept inside a refrigerator

indoors gave the lowest value of 38 r." Thus, although References 5 and 10 agree on the maximum and minimum readings, they differ on the range of doses to personnel.

Dr. Payne S. Harris of Los Alamos National Laboratory (then LASL) listed film badge numbers and their respective doses (Reference 9). These data, in Table 2, approximate those film badge readings quoted in References 1 and 2, with a range in personnel dose of 40-52 R and maximum and minimum doses of 98 R and 37.5 R, respectively, depending on whether the badge remained outside or inside for the exposure period. For comparison with calculated doses, the Reference 9 doses are used.

Table 2. Film badge dosimetry results,
Rongerik Atoll, 1-2 March 1954.

<u>Badge#</u>	<u>Location</u>	<u>Dose (R)</u>
309	Refrigerator	37.5
310	CN 408	40.0
311	CN 416	40.0
312	CN 411	44.0
313	CN 419	52.0
314	On tent pole	98.0
315	On barracks	82.0
316	Refrigerator	38.0
317	Refrigerator	38.5
318	Refrigerator	37.5
319	Refrigerator	37.5
320	Refrigerator	37.5

3.3 COMPARISON OF ESTIMATED FILM BADGE DOSES WITH FILM BADGE DATA.

All film badges except badge # 313 were turned in and placed in the mess hall at approximately 1130 hours, 2 March (H+28.75), where they remained until evacuation at 1800 hours (H+35.25). Therefore, the estimated film badge doses in Table 1, which are based on personnel movements on the island until evacuation, cannot be compared directly with those film badge readings in Table 2. In order to make this comparison

for a specific badge, the exposure of the film badge during the period it was in the mess hall (H+28.75 to H+35.25) must be added to the calculated film badge dose as of H+28.75 hours. This "mess hall dose" is approximately 12 rem.

From the personnel interviews it is possible to calculate the H+28.75 film badge dose for two of the film badges whose readings are given in Table 2 (badge #311 and #312). In addition, it is possible to calculate the H+28.75 hour dose for the badge that was hung on the tent post in the Army area (badge #314). Table 3 compares the calculated film badge dose at H+35.25 hours (the calculated dose at H+28.75 hours plus the "mess hall dose") and the corresponding film badge reading (from Table 2) for these three film badges. The activity scenarios or protection factors pertinent to the other badges are insufficiently known to permit meaningful comparisons.

Table 3. Comparison of calculated film badge doses with dosimetry.

<u>Film Badge</u>	<u>CN</u>	<u>Calculated Dose @H+28.75 hrs</u>	<u>Calculated Dose @H+35.25 hrs</u>	<u>Film Badge Reading (Table 2)</u>	<u>Ratio</u>
311	416	32 rem	44 rem	40 rem	1.10
312	411	38	50	44	1.14
314	N/A	100	112	98	1.14

From Table 3, the calculated dose to a specific film badge at H+35.25 hours is approximately 10-14 percent higher than the actual film badge reading. That these ratios are similar under varying circumstances of exposure suggests that there is a systematic overestimation of dose in the foregoing analysis. The difference between the calculated doses and the film badge readings could easily be accounted for in the uncertainties associated with defining the free-field radiation environment and protection factors on Rongerik. Thus, the estimated doses in Table 1 are reduced by a factor of 1.1 in order to be consistent with the dosimetry results. This adjustment in the Table 1 doses yields the best estimates of film badge doses received by the personnel on Rongerik. Such an approach provides the most consistency when the exposures are characterized by a mix of film badge readings and calculated doses. Results are presented in Section 5.

SECTION 4

TECHNICAL APPROACH — INTERNAL DOSE

The technical approach used to estimate the organ-specific radiation dose commitments (i.e., committed dose equivalent) for the American military personnel exposed on Rongerik Atoll is based on the following data:

- o Radionuclide inventory data for Shot Bravo calculated by Lawrence Livermore National Laboratory.
- o Urinalysis data for the exposed Americans obtained by Los Alamos National Laboratory.

The basic approach used to calculate the dose commitments has four steps.

- o The calculated radionuclide inventory provides the time-dependent relative activity of the radionuclides present in the fallout.
- o The inventory is normalized to the actual activity intake by relying on post-shot urinalysis to determine the amount of I-131 activity intake.
- o The internal dose pathways are examined to determine that doses are high-sided through use of large-particle inhalation dose conversion factors (rem/ μ Ci), which are developed by modifying published ingestion dose conversion factors.
- o Organ-specific dose commitments (rem) are calculated by combining the normalized activity inventory data with the large particle inhalation dose conversion factors.

A description of each of the above steps is provided in the subsections below.

4.1 RELATIVE ACTIVITY.

The basic fallout radiological data calculated for Shot Bravo was provided by Lawrence Livermore National Laboratory (LLNL) (Reference 11). The calculations were performed using the isotope generation and depletion code ORIGEN (Reference 12). Modifications to the ORIGEN-calculated results were made based on radiation chemistry data for Bravo available to LLNL. The LLNL-calculated data for Bravo thus reflect the device-specific characteristics (such as fissile nuclides, neutron energy, and light/heavy element production) that are necessary to correctly specify the radionuclide content of the fallout material. The unfractionated inventory is used because it high-sides organ doses derived from an iodine-based bioassay and because no specific level of fractionation can be substantiated. Plutonium nuclide information does not explicitly appear in Reference 11, but has been derived based on guidance from the author.

4.2 INVENTORY NORMALIZATION.

As previously mentioned, urine samples were collected from the exposed American servicemen and sent to US laboratories for analysis. These afford the opportunity for a more accurate dose calculation than possible from first-principle physical considerations. On-site radiation surveys were late (D+9 days) and did not facilitate an internal exposure analysis (Reference 8). The urinalysis data collected by Dr. Payne S. Harris of the Los Alamos National Laboratory are the most useful for this dose calculation because they are based on a comparatively early (D+17 days) sample collection and are well documented (Reference 9). The urinalysis results reported by the Naval Radiological Defense Laboratory involved samples collected much later (D+44 days and after), for which the available documentation is weak (Reference 13). It is understood that urinalysis results were also reported by medical personnel at Tripler Army Hospital; however, apparently the sample collection times were so late that meaningful results were not reported (Reference 4).

The urinalysis data reported by Dr. Harris for the American servicemen were based on a 15-min, pooled urine sample collected on 18 March 1954. The urine

samples were analyzed for plutonium, calcium-45, strontium-89, ruthenium-103, iodine-131, and barium-140. The I-131 analyses of the pooled urine samples of the servicemen and Marshallese provided the basis for thyroid dose estimates by Dr. Harris and later researchers. Dr. Harris judged that I-131 analysis was the most accurate; it also led to the highest doses to the servicemen (Reference 9).

The basic finding of the I-131 analysis of the pooled urine sample was the determination that, for an average individual, the amount of I-131 excreted in urine over a 24-hour period at the time of the urine sample collection (D+17 days) was 4.0 microcuries. That result is used to determine the amount of I-131 intake.

The amount of a radionuclide excreted in a 24-hour urine sample is related to the amount of the radionuclide initially intaken (that is, inhaled and/or ingested) by

$$A_u(t) = Q_I \cdot f_I \cdot Y(t) \cdot F_u \quad (5)$$

where

$A_u(t)$ = the amount (μCi) of the radionuclide in a 24-hour urine sample taken at time t ,

Q_I = the amount (μCi) of the radionuclide intake,

f_I = the fractional amount of the radionuclide intake that is transferred to the blood,

$Y(t)$ = the ratio of the amount (μCi) of the radionuclide excreted with a 24-hour period to the amount (μCi) of the radionuclide uptaken by the blood, and

F_u = the fraction of the total daily excretion that is present in the urine.

To solve this equation, not only must the time t of excretion relative to that of intake be known, but also, for a radionuclide mixture, the time after detonation/release. In

addition, the pathway may influence the body absorption fraction f_1 and $Y(t)$ may be influenced by the presence of parent radionuclides.

As mentioned above, the value of $A_u(D+17)$ is 4.0 nCi or 0.004 μCi for I-131. For iodine, the applicable biological parameters are $f_1 = 1.0$ and $F_u = .86$ (derived from References 14, 15, and 23). The value of $Y(t)$ is derived from Reference 16, which contains a recent retention model. This model does not differentiate between the relatively short time from intake to organ uptake, which is satisfactory due to the rapid absorption of iodine by the body. The value of $Y(D+17)$ used is for the first day (effectively instantaneous relative to 17 days) intake: $Y(D+17) = 2.87 \times 10^{-4}$ $\mu\text{Ci/day}$ per μCi uptake. This value reflects excreted I-131 that entered the body as I-131. However, at an early (shot-day) time of intake, the parent radionuclide Te-131m (half-life of 30 hours) exists in modest abundance--about 10 percent as much as I-131, by mass (Reference 11). Although tellurium is only partially absorbed into the body, at least half of the Te-131m decays prior to elimination from the body, and the iodine produced is fully absorbed. Therefore, up to about 7 percent of the excreted I-131 on D+17 could be from intake of Te-131m. With this adjustment, $Y(D+17) = 3.07 \times 10^{-4}$.

Using the values of $A_u(D+17)$, f_1 , $Y(D+17)$, and F_u cited above, the amount of the I-131 intake, Q_I , is 15 μCi . This value of Q_I is used to normalize the radionuclide inventory. Based on I-131 representing 0.2-0.3 percent of the total fission product activity (Reference 11) at the estimated time of intake (discussed subsequently), the total activity intake of fission products was about 5 to 7 mCi.

4.3 PATHWAYS AND DOSE CONVERSION FACTORS.

Researchers have arrived at various conclusions as to the dominant pathways for exposure to the Bravo contamination. The assumption of an inhalation pathway was made by Cole based on a comparison of the internal and external radiation dose estimates for the Americans on Rongerik and the natives on Rongelap (Reference 4). Cole's basic logic was that since the ratio of the internal and external doses was practically the same for both the Americans and the natives, the exposure pathways must have been similar (i.e., inhalation) and not related to the personal hygiene or

consumption habits that were dissimilar. However, the latest determination of internal dose for the Marshallese (Reference 19) considers ingestion to be the dominant pathway and arrives at significantly greater thyroid doses than previously calculated. Dr. Harris considers that ingestion through food or drinking water also contributed to the internal dose of the servicemen on Rongerik (Reference 9).

Scoping calculations suggest that both inhalation and ingestion pathways could have contributed significantly to internal dose at Rongerik, but that inhalation could not have dominated. As derived in Reference 20, the inhaled activity (curies) from full exposure to descending fallout is roughly 10^{-5} x peak gamma intensity (R/hr) x time of fallout after detonation (hr). The values of these parameters obtained in Sections 2 and 3 imply an inhaled activity on the order of 1 mCi. Ingestion could have resulted in virtually all of the activity intake, if, for example, food on a plate exposed to 15 minutes of significant fallout deposition were entirely ingested. At the time of the peak intensity, the areal concentration of activity exceeded 1 Ci/m^2 (Reference 11). With a plate size of 0.1 m^2 and exposure for 5 percent of the estimated duration of significant fallout, an intake of 5 mCi is implied.

Later pathways are less likely to have contributed significantly to internal dose. Fallout, which had accumulated to a noticeable depth, would have been brushed or washed off of objects pertinent to ingestibles; at lower levels of contamination, an unrealistically large surface area would have to be involved for ingestion at the mCi level. Inhalation during the following day of contaminants resuspended by wind or personal agitation would have been minor.

The particle-size characteristics of the airborne fallout material are adequately known. Based on considerations of particle fall rate, Bravo cloud height, and time after detonation, it is estimated that the deposited fallout particles were in the size range of 55 to 120 μm diameter. Analysis of soil samples on Rongerik indicated that most of the radioactivity in soil was associated with particles in the size range of 60 to 200 μm diameter (Reference 13). In the context of aerosols, the deposited material consisted of large particles (greater than 10 μm in diameter). For dosimetric purposes, differentiation among large particle sizes is unnecessary, as shown below.

For most typical radiological contamination situations involving the inhalation of radioactive material, organ dose commitments are calculated using published inhalation dose conversion factors. Inhalation dose conversion factors are available for activity on particles of less than $10\mu\text{m}$ in diameter; however, inhalation dose conversion factors for large particles are not available in the published literature. Accordingly, for the purpose of this dose calculation a set of large-particle inhalation dose conversion factors is developed.

For particles greater than $10\mu\text{m}$ diameter, essentially all of the inhaled material is initially deposited in the nasal or nasopharyngeal region of the respiratory tract (Reference 17). This material is removed by chemical (absorption) and mechanical processes, and goes either to the blood, with subsequent transfer to other body organs, or to the gastrointestinal (GI) tract, with subsequent transfer to the blood and further transfer to other body organs or bodily elimination by excretion. The "pulmonary clearance classification" of the inhaled material (roughly analogous to the solubility of the material) determines the amount of the material that is initially transferred to the blood or the GI tract; for example, 99 percent of an "insoluble" material will go to the the GI tract with only 1 percent going to the blood, whereas a "very soluble" material will be divided evenly between the GI tract and the blood (Reference 16).

As ingestion plays a major role in the fate of an inhaled large particle, the large-particle inhalation dose conversion factors used for the present dose calculation are obtained by modifying ingestion dose conversion factors. This modification is necessary to properly account for that portion of the inhaled material that is initially transferred to the blood, rather than to the GI tract. The metabolic data needed to construct large-particle inhalation dose conversion factors are given in ORNL/NUREG/TM-190 (Reference 16) for the most radiologically significant radionuclides. The TM-190 dose conversion factors, based on the most current dosimetry information, are used in conjunction with these. For those radionuclides not addressed in TM-190, unmodified ingestion dose conversion factors are taken from NUREG-6172 (Reference 18).

Large-particle inhalation dose conversion factors (where derivable) are used to high-side estimates of internal dose, despite the apparent dominance of the ingestion pathway. For some radionuclides, the original deposition of inhaled particles in the nasopharyngeal region affords a significantly greater absorption into the body than occurs in the GI tract. A minor contribution to GI tract doses is depleted through this pathway, but lung dose is greatly increased. For iodine, absorption is essentially complete with either pathway. Thus, the iodine-dominated thyroid dose is insensitive to the mix of ingestion and large-particle inhalation contributions to the total activity intake (so long as these occur at about the same time). Moreover, the use of the calculated I-131 intake to normalize the radionuclide inventory is independent of this mix.

Dose calculations are made for intake at 9 hours after detonation. In the early portion of significant fallout deposition, this high-sides organ doses by including greater activities of fast-decaying radionuclides. The large-particle dose conversion factors used in the calculation are listed in Reference 21.

4.4 ORGAN-SPECIFIC DOSE COMMITMENTS.

The 50-year dose commitment to organ j , D_j , resulting from the intake of a mixture of radionuclides is given by

$$D_j = \sum_i Q_I^i \cdot DCF_i^j \quad (6)$$

Q_I^i , the amount of intake of radionuclide i , is determined from the radionuclide inventory as normalized by the I-131 activity intake developed from the urinalysis data. The dose conversion factor, DCF_i^j , for organ j due to the intake of radionuclide i is as discussed previously.

SECTION 5

RESULTS AND COMPARATIVE ANALYSIS

5.1 RESULTS.

EXTERNAL DOSE

The gamma radiation dose to the servicemen on Rongerik Atoll is reconstructed as follows. The radiological environment resulting from Shot Bravo fallout is determined from time-of-arrival data on Rongerik, modeled deposition, measured decay rates, and a late reading on Rongerik. Personnel activity scenarios are considered in conjunction with radiation shielding factors to arrive at film badge doses on an individual basis. Although film badge readings are available, they inadequately relate to personnel dose. Readings that reflect well-specified exposures indicate good agreement with corresponding calculated doses. A small, but systematic, difference indicates an adjustment to the calculated personnel doses. Average doses to the identifiable subgroups of Rongerik personnel are as follows:

<u>Group</u>	<u>Number of People</u>	<u>Average Dose (rem)</u>
First Evacuation	8	33 ⁺² ₋₁
Second Evacuation		
Air Force	17	38 ⁺³ ₋₄
Army	3	43 ⁺³ ₋₂

Reconstructed doses are presented by case number in Table 4, with individual scenario information included as available.

INTERNAL DOSE

Organ-specific 50-year dose commitments resulting from the large-particle inhalation and ingestion of Bravo fallout radionuclides are given in Table 5. The

Table 4. Reconstructed gamma radiation doses for military personnel stationed on Rongerik Atoll, 1 and 2 March 1954.

<u>Air Force</u>	<u>Army</u>	<u>Clinical Number Assigned</u>	<u>Film Badge Number* (Reading)</u>	<u>Reconstructed Dose (rem)</u>
FIRST EVACUATION				
X		401		33**
X		402		33**
X		403		33**
X		404		35
X		405		33**
X		406		32
X		407		32
X		408	#00310 (40R)	33**
SECOND EVACUATION				
X		409		41
X		410		38**
X		411	#00312 (44R)	41
X		412		34
X		413		38**
	X	414		46
X		415		39
X		416	#00311 (40R)	39
X		417		40
X		418		37
X		419	#00313 (52R)	52***
X		420		38
X		421		38**
X		422		36
	X	423		43
X		424		38**
	X	425		43
X		426		34
X		427		40
X		428		38**

*Reference 9

**Mean dose for group

***Actual film badge reading is given because it represents entire exposure.

Table 5. Fifty-year internal dose commitments for military personnel stationed on Rongerik Atoll, 1 and 2 March 1954.

<u>Organ</u>	<u>Dose Commitment (rem)</u>
Thyroid	195
Lower large intestine wall	76
Upper large intestine wall	44
Small intestine wall	13
Lung	13
Stomach wall	6.4
Endosteum (bone surfaces)	3.4
Liver	1.6
Bone	1.4
Red marrow	1.3
Kidney	1.2
Pancreas	0.9
Testes	0.6

calculation of internal dose commitments is based on an intake at 9 hours after Shot Bravo, with large-particle dose conversion factors applied to the radionuclide inventory to that time. The inventory, which is based on radiochemical analysis of Bravo, is normalized to the activity level of iodine-131. Intake of this radionuclide is determined from the body metabolism of iodine that led to a measured activity of I-131 in urine samples taken 17 days after Bravo.

The calculated thyroid dose commitment of 190 rem is by far the largest organ dose commitment, and the dose is accrued comparatively rapidly. Short-lived radionuclides dominate the thyroid dose, which is seven times that from I-131 alone. If the intake had been at 13 hours after Bravo, the thyroid dose would be six times that from I-131 (based on the Reference 11 radionuclide inventory). Thus, the thyroid dose is not sensitive to actual time of intake within the fallout deposition period.

The only other dose commitments that exceed the dose from external gamma radiation are for the sections of the large intestine. Actinide emitters, produced by the neutron activation of U-238 during the detonation, are significant to these doses. Because GI tract doses are dominated by the ingested contents in transit rather than by absorbed radionuclides, these doses are also accrued rapidly.

Caution is advised in interpreting these calculated internal doses for specific individuals. Based on a pooled urine sample, the calculated doses represent an average or nominal dose estimate and do not necessarily pertain to a specific individual. However, limited individual urine sampling reported by Dr. Harris (Reference 9) indicated only a modest spread of results. The counts of the individual samples were consistent to within 40 percent of their mean. Also, the implied intake of I-131 agreed well with that from the pooled sample.

5.2 COMPARISON WITH PREVIOUS DOSE DETERMINATIONS.

Dose determinations for personnel on atolls downwind of CASTLE Bravo have been principally of two types: the initial estimates developed in the mid-1950s, shortly after the event, and the long-term, continuing followup by Brookhaven

National Laboratory on the affected Marshallese. While the servicemen on Rongerik Atoll do not fall under the Brookhaven charter, they do obtain limited mention in Brookhaven reports with respect to the consistency of external dose estimates. Moreover, the internal dose methodologies for the two groups are perforce similar, as the underlying data are related. Consequently, as the current Brookhaven report (Reference 19) provides a synthesis of earlier work, it affords the best available comparison with this analysis.

EXTERNAL DOSE

Previous studies have the following features in common for dose estimates to the servicemen on Rongerik: doses are based on free-field intensities, are reported as air exposure (R), and thus are non-specific as to individual's activities. These estimates have recently been referred to in terms of rad or rem, which further blurs the distinction between them and the calculated film badge doses that are presented in this report. Thus, the previous estimates, which are generally in the range of 70-100 R, would correspond to about 50-70 rem with the use of the 0.7 rem/R film badge conversion factor. With the inclusion of the protection factor of about 2 for time indoors, the estimates would be reduced further and adequately correspond to the present results. Earlier reports displayed no interest in arriving at individualized estimates; it is not even clear that the 1950's researchers had access to the full range of then-classified information that would have been required for such estimates.

Reference 19 lists along with its current estimates of whole body external doses to the Marshallese a corresponding value for Rongerik, 81 rad. While this value is useful for comparing doses for various Marshallese situations, it does not apply to the servicemen. The value was treated as if it were the whole body dose and thus additive, without adjustment, to an internal dose in rem. This was done for the Marshallese to arrive at total thyroid doses. The 81 rad was arrived at by application of a Marshallese living-style factor to an estimate of integrated intensity in free air. No consideration was given to the substantially greater shielding of the Rongerik servicemen (Reference 22).

One of the first estimates of Rongerik external whole-body dose is cited in Reference 19 as 78 rad, from WT-939 (Reference 5). This value came through the use of a secondary reference (Reference 22); WT-939 actually provides a range of possible air doses (R), from which it selected a value of 86 R. While WT-939 did report final results in terms of R, it contains a discussion of corresponding depth doses in tissue. The magnitude of depth doses, relative to the air dose, are consistent with the aim of a film badge reading to adequately reflect depth dose. Correspondingly lower film badge doses would follow from these relationships.

Even after conversion to depth dose/film badge dose, the WT-939 dose estimate remains greater than the present value for two reasons: how it considered shielding and how it arrived at an integrated intensity in air. Without the benefit of WT-938 (Reference 3), published later, WT-939 incorporated no specifics on personnel movement; an estimate was used of one-half of the time outdoors with no protection and one-half of the time indoors with a protection factor of 2. Based on the personnel scenarios in WT-938, however, it is deduced that a considerably greater fraction of the time was spent indoors, and more of that was in buildings with a $PF > 2$ than $PF < 2$. In fact, for the lowest dose case (CN406), detailed in the Appendix, the time-averaged PF was greater than 2. Thus, the calculated film badge dose for this case is less than two-thirds of what WT-939 would suggest -- $0.65 \times 0.7 \times 86 = 39$ rem, close to the present value of 32 rem.

The WT-939 estimate of integrated intensity in air relies on an assumed duration of fallout deposition. While the present analysis initially follows this approach, the results are adjusted for actual film badge readings. Both indoor and outdoor badges, when shielding and duration of exposure are properly accounted for, imply a consistent adjustment factor. Lacking any known systematic error in the film badge readings, the normalization to badge readings provides the most credible personnel dose determination. The implied integrated intensity in air is about 15 percent less than in WT-939. This resolves the remnant difference in personnel dose estimates.

No other known estimate of Rongerik external dose is cited in Reference 19. A computerized theoretical treatment of Bravo fallout deposition leads to dose estimates that differ with other calculations for all atolls studied. The wind data available for this approach are too sparse, leading to results that are not credible (Reference 22).

INTERNAL DOSE

The current Brookhaven report has a number of features that could make it useful for comparison with internal dose estimates for the Rongerik servicemen: closely related urinalysis data for the Marshallese and the servicemen are utilized in similar internal dose methodologies; the latest thyroid dose estimates (to the Marshallese) exceed any of the referenced previous estimates; and use is made of ICRP-30 (Reference 14), a likely foundation for the present analysis had all of the metabolic and dosimetric information for specific radionuclides been published when this work was initiated.

Both the Brookhaven and the present assessments utilize I-131 counts from urinalyses to arrive at initial body burdens of I-131. From this, Brookhaven determines the levels of other iodine isotopes that lead to thyroid dose. The present analysis goes beyond, with use of Reference 11, to determine the levels of radionuclides generally, and from them and References 16-18, the doses to all relevant organs. However, the excretion values used in Reference 19 do not specifically relate to this analysis because of the longer duration of intake, the earlier urine sampling time, and the non-standard daily urine sample size, which is considered to influence iodine metabolism.

The common basis for the ORNL-TM-190 (Reference 16) and ICRP-30 (References 14, 15) treatments of iodine metabolism is Reference 23. TM-190 has applied the Reference 23 prescription numerically as a function of time for various body compartments; ICRP-30 gives approximated time constants for these compartments, from which the user can develop functional relationships. When this is done, the agreement between I-131 body burdens deduced from ICRP-30 and TM-190 is good, especially with the approximations taken into account. However, this agreement exists only for the more recent version of iodine metabolism expressed in ICRP-30, Part 3 (Reference 15) -- ICRP-30, Part 1 (Reference 14) has since been acknowledged as incorrectly stating the iodine parameters (References 15, 24). The most recent work on radionuclide metabolism, Reference 25, tabulates daily excretion fractions that are in close accord with this analysis.

5.3 TOTAL DOSE.

The total dose to each organ is the sum of the external and internal dose contributions. For consideration of risk assessment, the long-term aspect of certain dose commitments must be included. However, the important contributions of internal dose to the total dose of the Rongerik personnel were only for organs in which the internal dose is rapidly accrued. Approximate total values are 230 rem to the thyroid, 115 rem to the lower large intestine wall, 85 rem to the upper large intestine wall, and about 40 to 50 rem to all other organs.

Uncertainties in the dose calculations are not readily determined. For external dose, they are judged to be small in that the calculations rely substantially on film badge readings. No information is available as to the accuracy of this dosimetry. For internal dose, the referenced metabolic information lacks estimates of uncertainty. Throughout the calculations, a high-sided estimate is made when a choice of specific parameters is available.

SECTION 6
LIST OF REFERENCES

1. The Effects of Nuclear Weapons, S. Glasstone and P.J. Dolan, US Department of Defense and US Department of Energy, Washington, DC, GPO, 1976.
2. "Environmental Radioactivity," M. Eisenbud, Second Edition, Academic Press, New York, 1973.
3. "Exposure of Marshall Islanders and American Military Personnel to Fallout," WT-938-EX, R. Sharp and W.H. Chapman, Naval Medical Research Institute, Operation CASTLE, February 1959. AD/A 995 077.*
4. "Inhalation of Radioiodine from Fallout: Hazards and Countermeasures," R. Cole, Defense Civil Preparedness Agency, August 1972, AD 761 638.*
5. "Physical Factors and Dosimetry in the Marshall Island Radiation Exposure," WT-939, Addendum Report, Operation CASTLE, Project 4.1, C.A. Sondhaus and V.P. Bond, Naval Radiological Defense Laboratory, December 1955, AD 338 337.*
6. "Distribution and Intensity of Fallout," Operation CASTLE, WT-915, R.L. Steton, Naval Radiological Defense Laboratory, January 1956, AD 361 836.*
7. Analysis of Fallout Data, Part II, "Decay Characteristics of Radioactive Fallout," C.F. Miller, Naval Radiological Defense Laboratory-TR-221, 1958.
8. Letter Report, Subject: Radiological Survey of Downwind Atolls Contaminated by BRAVO, Herbert Scozillo, Jr., Hq. Task Unit 13, Task Group 7.1, TU-13-54-374, 12 March 1954.
9. Private Notes and Communications, Dr. P.S. Harris to SAIC Staff, 1982-1983.

* Available from NTIS; or for number appears before the asterisk.

** Available at OIC.

10. "Study of Response of Human Beings Accidentally Exposed to Significant Fallout Radiation," Operation CASTLE, WT-923, Project 4.1, E.P. Cronkite et al., Naval Medical Research Institute, January 1954, AD 465 295.*
11. "Results of Calculations of External Gamma Radiation Exposure Rates from Local Fallout and the Related Radionuclide Compositions of Selected US Pacific Events," UCRL-53555, H.G. Hicks, Lawrence Livermore National Laboratory, 1984, (A15) DE 84 010 678.*
12. "ORIGEN - The ORNL Isotope Generation and Depletion Code," ORNL-4628, M.J. Bell, Oak Ridge National Laboratory, May 1973.
13. "Nature and Extent of Internal Radioactive Contamination of Human Beings, Plants and Animals Exposed to Fallout," WT-936, Operation CASTLE, S.H. Cohn et al., Naval Radiological Defense Laboratory and Naval Medical Research Institute, 1957, AD 617 134.*
14. "Limits for Intakes of Radionuclides by Workers," International Commission on Radiological Protection, ICRP Publication 30, Part 1, Pergamon Press, New York, 1978.
15. _____, ICRP Publication 30, Part 3, Pergamon Press, New York, 1981.
16. Estimates of Internal Dose Equivalent to 22 Target Organs for Radionuclides Occurring in Routine Releases from Nuclear Fuel-Cycle Facilities, 2 vols., G.G. Killough et al., ORNL/NUREG/TM-190, Jun 78 and Nov 79, NUREG/CR-0150* and NUREG/CR-0150-V-2.*
17. Task Group on Lung Dynamics, "Deposition and Retention Models for Internal Dosimetry of the Human Respiratory Tract," Health Physics, 12(1966), p. 173-267.

*Available from NTIS; order number appears before the asterisk.

**Available at CIC.

18. "Age Specific Radiation Dose Commitment Factors for a One Year Chronic Intake," NUREG-0172 (PB-272-348), G.R. Hoenes and J.K. Soldat, Battelle Pacific Northwest Laboratories, November 1977.*
19. "Thyroid Absorbed Dose for People at Rongelap, Utirik, and Sifo on March 1, 1954," BNL 51882 *E. Lessard et al., Brookhaven National Laboratory, March 1985.
20. "Low Level Internal Dose Screen, CONUS Tests," DNA-TR-85-317, Defense Nuclear Agency, 22 December 1986.
21. "FIIDOS--A Computer Code for the Computation of Fallout Inhalation and Ingestion Dose to Organs," DNA-TR-84-375, Defense Nuclear Agency, December 1985.
22. Discussion with E. Lessard, Brookhaven National Laboratory, 25 March 1986.
23. "Quantitative Aspects of Iodine Metabolism in Man," D.S. Riggs, Pharmacological Reviews, Vol. 4 (1952), p. 284-370.
24. Discussion with K. Eckerman, member ICRP Committee 2, 11 March 1986.
25. "Interpretation of Bioassay Measurements," NUREG/CR-4884 (BNL 52063), E. Lessard, X. Yihua, K. Skrabale, G. Chabot, C. French, T. Lebone, J. Johnson, D. Fisher, R. Belanger, J. Lipsztein, Brookhaven National Laboratory, March 1987.

*Available from NTIS.

**Available at CIC.

APPENDIX

SAMPLE CASE INTERVIEW

After being evacuated to Kwajalein on 2 March 1954, all of the military personnel assigned to Rongerik Atoll were interviewed to determine their specific activities during the period 1 March until their subsequent evacuation on 2 March. Some of the interviews provided sufficient detail to determine an individual's activities, hence his whereabouts, on an hour-to-hour basis. Because the film badge dose estimates in this memorandum are based on seventeen such interviews, this appendix shows how these data are utilized in calculating the film badge doses.

Reference 3 contains a summary of the interviews obtained from the 28 military personnel after being evacuated to Kwajalein. The results of an interview with CN 406, extracted from Reference 3, is as follows:

CN 406

Duty. Rawinsonde Operator.

Activities. 1 March: On duty day of 1 March until 1630. Took shower before chow. Attended movie in mess hall.

2 March: Worked outside, helping to wash outside of mess hall.

Clothing. Wore shorts until 1530. Changed to long clothing.

Food and Drink. Ate evening meal 1 March. Had coffee for breakfast, 2 March. No lunch.

Fallout. Observed fallout at about 1500. "White ashes; light, floated down."

Film Badge. None. CN408 wore badge for group.

Evacuated. In first group (1245 hours, 2 March).

Note: High school education. Thirteen months remaining of present enlistment.

Based on the interview with CN 406, it is possible to estimate where this individual was at a particular time and, hence, determine what radiation protection was provided by the particular building he was in. When a particular structure was not specified, it was assumed he was outdoors (PF=1.0). Table 6 details the activities of CN 406 as inferred from his interview, and lists the calculated film badge dose for this individual during his stay on Rongerik.

Table 6. Calculated film badge dose for one airman (CN 406).

<u>1-2 MARCH 1954</u> <u>TIME (LOCAL)</u>	<u>TIME AFTER</u> <u>BURST (HRS)</u>	<u>ACTIVITY</u>	<u>PF</u>	<u>CUMULATIVE</u> <u>DOSE (rem)</u>
1430-1630	7.75-9.75	Radiosonde bldg-on duty	2.5	0.3
1630-1700	9.75-10.25	Shower	1.0	0.9
1700-1800	10.25-11.25	Mess hall-chow	2.0	2.0
1800-2300	11.25-16.25	Mess hall-movies	2.0	13.0
2300-0700	16.25-24.25	Barracks	2.9	23.6
0700-0800	24.25-25.25	Mess hall-chow	2.0	25.2
0800-1000	25.25-27.25	Washing down mess hall	1.0	31.3
1000-1245	27.25-30.0	Mess hall until evacuation	2.0	35.2*

*This total is subsequently reduced by 10 percent as described in Section 3.3.

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE

ARMED FORCES RADIOBIOLOGY RSCH INST
ATTN: DIRECTOR
ATTN: TECHNICAL LIBRARY

ASSISTANT SECRETARY OF DEFENSE
PUBLIC AFFAIRS
ATTN: ASD(PA)

ASSISTANT SECRETARY OF DEFENSE
ATTN: ASD(HA)

DEFENSE INTELLIGENCE AGENCY
2 CYS ATTN: RTS 2B

DEFENSE NUCLEAR AGENCY
ATTN: GC
ATTN: PAO
10 CYS ATTN: RARP
5 CYS ATTN: TITL

DEFENSE TECHNICAL INFO CENTER
12 CYS ATTN: DD

DEP UNDER SEC OF DEF FOR RSCH & ENGRG
ATTN: DUSDRE (RSCH & ADV TECH)

FIELD COMMAND DEFENSE NUCLEAR AGENCY
ATTN: FCL
2 CYS ATTN: FCLS
ATTN: FCTXE
ATTN: FCTXE
ATTN: FTTD W SUMMA

LAWRENCE LIVERMORE NATIONAL LAB
ATTN: DNA-LL

DEPARTMENT OF THE ARMY

HARRY DIAMOND LABORATORIES
ATTN: SLCIS-IM-TL (81100) (TECH LIB)

U S ARMY BALLISTIC RESEARCH LAB
ATTN: DRDAR-BLV-R J MALONEY

U S ARMY CTR OF MIL HISTORY
ATTN: LIBRARY

U S ARMY MED RSCH & DEV CMD
ATTN: SGRD:SD

U S ARMY NUC & CHEM AGENCY
ATTN: MONA-ZB(C DAVIDSON)

WALTER REED ARMY MEDICAL CENTER
ATTN: LIBRARY

DEPARTMENT OF THE NAVY

MARINE CORPS HISTORY & MUSEUMS
ATTN: HISTORICAL DIVISION

NATIONAL NAVAL MEDICAL CENTER
ATTN: DEPT OF RADIOLOGY
ATTN: MEDICAL LIBRARY

NAVAL HISTORICAL CENTER
ATTN: DD ALLARD

NAVAL MEDICAL COMMAND
ATTN: MEDCOM-21
ATTN: NM&S-00
ATTN: NM&S-09

NAVAL OCEAN SYSTEMS CENTER
ATTN: TECH LIB

NAVAL SEA SYSTEMS COMMAND
ATTN: SEA-08 M MILES

NAVAL WEAPONS EVAL FACILITY
ATTN: CLASSIFIED LIBRARY

OFC OF THE DEP CH OF NAVAL OPS
5 CYS ATTN: NOP 0455

DEPARTMENT OF THE AIR FORCE

AIR FORCE HISTORICAL RSCH CTR
ATTN: LIBRARY

AIR FORCE INSTITUTE OF TECH/EN
ATTN: LIBRARY/AFIT/LDEE

AIR FORCE WEAPONS LABORATORY
ATTN: SUL

AIR UNIVERSITY LIBRARY
ATTN: AUL-LSE

HQ USAF/SGPT
4 CYS ATTN: SGPT/COL GIBBONS

HUMAN SYSTEMS DIVISION (AFSC)
ATTN: LIBRARY SCL-4

USAF OCCUPATIONAL & ENV HEALTH LAB
4 CYS ATTN: AFNTPR
ATTN: R2

DEPARTMENT OF ENERGY

DEPARTMENT OF ENERGY
ATTN: OMA, DP-22

DEPARTMENT OF ENERGY
ATTN: B CHURCH, HPD

DEPARTMENT OF ENERGY
ATTN: J THIESEN
ATTN: TECHNICAL INFO CTR

LAWRENCE LIVERMORE NATIONAL LAB
ATTN: L-453 L ANSPAUGH
ATTN: L-53 TECH INFO DEPT LIB
ATTN: Y NG

LOS ALAMOS NATIONAL LABORATORY
ATTN: F601 T DOWLER
ATTN: ITO D STILLMAN
ATTN: P WHALEN

DNA-TR-86-120 (DL CONTINUED)

OAK RIDGE NATIONAL LABORATORY
ATTN: C RICHMOND
ATTN: G KERR

OAK RIDGE NATIONAL LABORATORY
ATTN: T JONES

REYNOLDS ELECT AND ENGR CO. INC
ATTN: CIC
ATTN: LST
ATTN: W BRADY

OTHER GOVERNMENT

CANCER CENTER, NIH
ATTN: A KNUDSON

CENTRAL INTELLIGENCE AGENCY
ATTN: OFFICE OF MED SERVICES

DEPARTMENT OF COMMERCE
ATTN: C KUYATT
ATTN: J HUBBELL
ATTN: M EHRLICH

DEPT OF HEALTH & HUMAN SERVICES
ATTN: OFC OF REGULATION REVIEW

DEPT OF HEALTH & HUMAN SVCS
ATTN: R MURPHY

DEPARTMENT OF LABOR
ATTN: S WEINER

DEPT OF HEALTH & HUMAN SERVICES
ATTN: C SILVERMAN
ATTN: G JOHNSON
ATTN: J VILLFORTH

ENVIRONMENTAL PROTECTION AGENCY
ATTN: P MAGNO
ATTN: T THORSLUND

ENVIRONMENTAL PROTECTION AGENCY
ATTN: N NELSON

FEDERAL EMERGENCY MGT AGENCY
ATTN: C SIEBENTRITT
ATTN: H TOVEY
ATTN: OFC OF RSCH/NP H TOVEY

LIBRARY OF CONGRESS
ATTN: SCIENCE & TECHNOLOGY DIV

NATIONAL CANCER INSTITUTE, NIH
ATTN: B WACHOLZ
ATTN: G BEEBE
ATTN: V ZEVE

NATIONAL CANCER INSTITUTE, NIH
ATTN: C LAND
ATTN: J FRAUMENI
ATTN: W BLOT

NATIONAL CANCER INSTITUTE, NIH
ATTN: J GART

NATIONAL CANCER INSTITUTE, NIH
ATTN: A RABSON
ATTN: D PISTENMAA
ATTN: J WYNGAARDEN

NATIONAL INSTITUTE FOR OCCUPATIONAL
ATTN: W MURRAY

NATIONAL LIBRARY OF MEDICINE, NIH
ATTN: LIBRARY

NATIONAL SCIENCE FOUNDATION
ATTN: P HARRLMAN

SUBCOMMITTEE OF NUC REGULATORY
ATTN: J CURTISS

U S HOUSE OF REPRESENTATIVES
ATTN: SUBCOMMITTEE ON MIL PER & COMP

U S HOUSE OF REPRESENTATIVES
ATTN: SUBCOMMITTEE ON HEALTH & ENVIR

U S HOUSE OF REPRESENTATIVES
2 CYS ATTN: MAJORITY COUNSEL

U S PUBLIC HEALTH SERVICE
ATTN: LIBRARY

U S SENATE
ATTN: J MCGOVERN

U S SENATE
3 CYS ATTN: MAJORITY COUNSEL

U S SENATE
ATTN: S ULM

VETERANS ADMIN MEDICAL CENTER
ATTN: K LEE

VETERANS ADMIN MEDICAL CENTER
ATTN: D MCGREGOR

VETERANS ADMIN MEDICAL CENTER
ATTN: C TESSMER

VETERANS ADMIN WADSWORTH HOSPITAL CTR
ATTN: T MAKINODAN

VETERANS ADMINISTRATION HOSPITAL
ATTN: R YALOW

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

[illegible]

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECCTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION-RO
ATTN: DIRECTOR

DNA-TR-86-120 (DL CONTINUED)

VETERANS ADMINISTRATION:RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION:RO
ATTN: DIRECTOR

VETERANS ADMINISTRATION:RP
ATTN: DIRECTOR

WHITE HOUSE (THE)
ATTN: OFC OF POLICY DEV(DP)

DEPARTMENT OF DEFENSE CONTRACTORS

BDM CORP
ATTN: J BRADDOCK

JAYCOR
6CYS ATTN: A NELSON, INFO SYS DIV

KAMAN SCIENCES CORPORATION
ATTN: DASIAC

LOUISIANA UNIV SCH OF MED, SHREVEPORT
ATTN: LIBRARY

NATIONAL ACADEMY OF SCIENCES
ATTN: C ROBINETTE

NEBRASKA UNIVERSITY OF
ATTN: LIBRARY

NORTHROP CORP
ATTN: Z SHANFIELD

OHIO STATE UNIVERSITY
ATTN: LIBRARY

PACIFIC SIERRA RESEARCH CORP
ATTN: H BRODE, CHAIRMAN SAGE

R & D ASSOCIATES
ATTN: C K B LEE

R & D ASSOCIATES
ATTN: A DEVERILL

RADIATION RSCH ASSOCIATES, INC
ATTN: N SCHAEFFER

RAND CORP
ATTN: P DAVIS
ATTN: TECH LIBRARY

RAND CORP
ATTN: B BENNETT

SCIENCE APPLICATIONS INTL CORP
2 CYS ATTN: C THOMAS
2 CYS ATTN: J GOETZ
2 CYS ATTN: J KLEMM
ATTN: J MCGAHAN
2 CYS ATTN: J PHILLIPS

SCIENTIFIC INFO SERVICES, INC
ATTN: LIBRARY

FOREIGN

CANADIAN EMBASSY
ATTN: LIBRARY

EDF RETN 1
ATTN: LIBRARY

INDIAN COUNCIL OF MEDICAL RSCH
ATTN: A TASKAR

MAURICE DELPLA
ATTN: M DELPLA

MCGILL UNIVERSITY
ATTN: R OSEASOHN

PRESIDENTE UMBERTO COLOMBO
ATTN: LIBRARY

PUERTO RICO SCH OF MED, UNIV OF
ATTN: LIBRARY

UNITED KINGDOM SCI MISSION
ATTN: MIL LIASION FOR DR RIDLEY

DIRECTORY OF OTHER

AKRON PUBLIC LIBRARY
ATTN: GOVT PUB LIBRARIAN

ALABAMA UNIVERSITY OF
ATTN: DIR OF LIBRARIES (REGIONAL)

ALASKA UNIVERSITY OF
ATTN: DIRECTOR OF LIBRARIES

ALLEN COUNTY PUBLIC LIBRARY
ATTN: LIBRARIAN

ANAHEIM PUBLIC LIBRARY
ATTN: LIBRARIAN

ARIZONA UNIVERSITY
ATTN: LIBRARIAN

ARKANSAS COLLEGE LIBRARY
ATTN: LIBRARY

ARKANSAS LIBRARY COMM
ATTN: LIBRARY

ARKANSAS UNIVERSITY OF
ATTN: GOVT DOCUMENTS DIV

BEMIOJI STATE COLLEGE
ATTN: LIBRARY

BRIGHAM YOUNG UNIVERSITY
ATTN: DOCUMENTS COLLECTION

BROOKHAVEN NATIONAL LAB
ATTN: A B BRILL
ATTN: E CRONKITE
ATTN: M BENDER
ATTN: TECHNICAL LIBRARY
ATTN: V BOND

BURLINGTON LIBRARY ATTN: LIBRARIAN	DETROIT PUBLIC LIBRARY ATTN: LIBRARIAN
CALIFORNIA INSTITUTE OF TECH ATTN: E LEWIS ATTN: R CHRISTY	DICKINSON STATE COLLEGE ATTN: LIBRARIAN
CALIF STATE LIBRARY (REGIONAL) ATTN: LIBRARIAN	DREW, UNIVERSITY OF ATTN: LIBRARY
CALIFORNIA STATE UNIVERSITY ATTN: LIBRARIAN	DULUTH PUBLIC LIBRARY ATTN: DOCUMENTS SECTION
CALIFORNIA, UNIVERSITY ATTN: GOVT DOCUMENTS DEPT	EASTERN BRANCH ATTN: LIBRARIAN
CALVIN T. RYAN LIBRARY ATTN: GOVT DOCUMENTS DEPT	EL PASO PUBLIC LIBRARY ATTN: DOCS & GENEALOGY DEPT
CENTRAL FLORIDA UNIV OF ATTN: LIBRARY DOCS DEPT	ENOCH PRATT FREE LIBRARY ATTN: DOCUMENTS OFFICE
CENTRAL WASHINGTON UNIVERSITY ATTN: LIBRARY DOCS SECTION	EVANSVILLE & VANDERBURGH CO PUB LIB ATTN: LIBRARIAN
CHARLESTON COUNTY LIBRARY ATTN: LIBRARIAN	FLORIDA STATE UNIVERSITY ATTN: DOCUMENTS DEPARTMENT
CHATTANOOGA HAMILTON CO ATTN: LIBRARIAN	FOND DU LAC PUBLIC LIB ATTN: LIBRARIAN
CHICAGO PUBLIC LIBRARY ATTN: GOVT PUBLICATIONS DEPT	FORT HAYS STATE UNIVERSITY ATTN: LIBRARIAN
CHICAGO, UNIVERSITY OF ATTN: P MEIER	FORT WORTH PUBLIC LIBRARY ATTN: LIBRARIAN
COLORADO STATE UNIV LIBS ATTN: LIBRARIAN	FRESNO COUNTY FREE LIBRARY ATTN: LIBRARIAN
COLORADO, UNIVERSITY OF ATTN: LIBRARY	GADSEN PUBLIC LIBRARY ATTN: LIBRARIAN
COLUMBIA UNIVERSITY ATTN: A BLOOM ATTN: LIBRARY	GEORGIA, MEDICAL COLLEGE OF ATTN: L STODDARD
COLUMBIA UNIVERSITY ATTN: DIV OF BIOSTATISTICS	GRAND RAPIDS PUBLIC LIBRARY ATTN: DIRECTOR OF LIBRARIES
COORDINATION & INFORMATION CTR ATTN: C/O REECD	GREENVILLE COUNTY LIBRARY ATTN: LIBRARIAN
CORNELL UNIVERSITY ATTN: W FEDERER	GUAM RFK MEMORIAL UNIV LIB ATTN: FED DEPOSITORY COLLECTION
DALLAS PUBLIC LIBRARY ATTN: LIBRARIAN	HARVARD SCHOOL OF PUB HEALTH ATTN: J BAILOR ATTN: LIBRARY ATTN: R REED
DAYTON & MONTGOMERY CITY PUB LIB ATTN: LIBRARIAN	HARVARD SCHOOL OF PUBLIC HEALTH ATTN: B MACMAHON
DECATUR PUBLIC LIBRARY ATTN: LIBRARIAN	HARVARD UNIVERSITY ATTN: W COCHRAN
DELTA STATE UNIVERSITY ATTN: LIBRARIAN	HAWAII, UNIVERSITY OF ATTN: Y MATSUMOTO

DNA-TR-86-120 (DL CONTINUED)

HOPKINSVILLE COMM COLL
ATTN: LIBRARIAN

IDAHO STATE UNIVERSITY LIBRARY
ATTN: DOCUMENTS DEPARTMENT

IDAHO, UNIVERSITY OF
ATTN: DIR OF LIB (REGIONAL)

ILLINOIS STATE LIBRARY (REGIONAL)
ATTN: GOVT DOCUMENTS BRANCH

INDIANA STATE LIBRARY (REGIONAL)
ATTN: SERIAL SECTION

INDIANA UNIVERSITY
ATTN: F PUTNAM

JOHNS HOPKINS UNIVERSITY
ATTN: A KIMBALL
ATTN: R SELTSER

KANSAS CITY PUBLIC LIBRARY
ATTN: DOCUMENTS DIV

KANSAS STATE LIBRARY
ATTN: LIBRARIAN

KANSAS STATE UNIV LIBRARY
ATTN: DOCUMENTS DEPT

KANSAS UNIV OF AGRI & APPLIED SCI
ATTN: H FRYER

KENTUCKY DEPT OF LIB & ARCHIVES
ATTN: DOCUMENTS SECTION

KENTUCKY, UNIVERSITY OF
ATTN: DIR OF LIB (REGIONAL)

KINGSTON HOSPITAL
ATTN: K JOHNSON

LOS ANGELES PUBLIC LIBRARY
ATTN: SERIALS DIV U S DOC

LOUISIANA STATE UNIVERSITY
ATTN: DIR OF LIB (REGIONAL)

MAINE UNIVERSITY, OF
ATTN: LIBRARIAN

MANKATO STATE COLLEGE
ATTN: GOVT PUBLICATIONS

MANTOR LIBRARY
ATTN: DIRECTOR OF LIBRARIES

MARATHON COUNTY PUBLIC LIBRARY
ATTN: LIBRARIAN

MARYLAND UNIVERSITY OF
ATTN: MCKELDIN LIBR DOCS DIV

MAUI PUBLIC LIBRARY
ATTN: LIBRARIAN

MEM HOSP FOR CANCER & ALLIED DISEASES
ATTN: P LIEBERMAN

MEM SLOAN-KETTERING CANCER CENTER
ATTN: J LAUGHLIN
ATTN: P MARKS

MEMPHIS SHELBY CO PUB LIB & INFO CTR
ATTN: LIBRARIAN

MERCER UNIVERSITY
ATTN: LIBRARIAN

MERCK, SHARP & DOHME INTL
ATTN: A BEARN

MESA COUNTY PUBLIC LIBRARY
ATTN: LIBRARIAN

MIAM PUBLIC LIBRARY
ATTN: DOCUMENTS DIVISION

MICHIGAN MED SCHOOL, UNIV OF
ATTN: J NEEL

MICHIGAN STATE LIBRARY
ATTN: LIBRARIAN

MICHIGAN, UNIVERSITY OF
ATTN: R CORNELL

MICHIGAN, UNIVERSITY OF
ATTN: F MOORE

MINNESOTA, UNIVERSITY OF
ATTN: J BEARMAN
ATTN: L SCHUMAN
ATTN: LIBRARY

MINOT STATE COLLEGE
ATTN: LIBRARIAN

MISSOURI, UNIVERSITY LIBRARY
ATTN: GOVERNMENT DOCUMENTS

MONTANA STATE LIBRARY
ATTN: LIBRARIAN

NASSAU LIBRARY SYSTEM
ATTN: LIBRARIAN

NATL COUNCIL ON RADIATION
ATTN: W SINCLAIR

NATRONA COUNTY PUBLIC LIBRARY
ATTN: LIBRARIAN

NEW HAMPSHIRE UNIVERSITY LIB
ATTN: LIBRARIAN

NEW MEXICO, UNIV OF
ATTN: C KEY
ATTN: R ANDERSON

NEW YORK PUBLIC LIBRARY
ATTN: LIBRARIAN

NEW YORK STATE LIBRARY
ATTN: DOCS CONT CULTURAL ED CTR

NEW YORK STATE UNIV OF
ATTN: LIBRARY DOCUMENTS SEC

NEW YORK UNIV MEDICAL CENTER
ATTN: N NELSON

NEW YORK UNIVERSITY
ATTN: A UPTON
ATTN: B POSTERNACK
ATTN: LIBRARY

NEWARK FREE LIBRARY
ATTN: LIBRARIAN

NORTH CAROLINA UNIV AT WILMINGTON
ATTN: LIBRARIAN

NORTH CAROLINA, UNIVERSITY OF
ATTN: LIBRARY FOR DEAN

NORTHERN ARIZONA UNIVERSITY LIB
ATTN: GOVERNMENT DOCUMENTS DEPT

NORTHERN IOWA UNIVERSITY
ATTN: LIBRARY

NORTHERN MICHIGAN UNIV
ATTN: DOCUMENTS

NORTHERN MONTANA COLLEGE LIBRARY
ATTN: LIBRARIAN

NORTH WESTERN MICHIGAN COLLEGE
ATTN: LIBRARIAN

NORTHWESTERN UNIVERSITY
ATTN: H CEMBER

OAK RIDGE ASSOCIATED UNIVERSITIES
ATTN: D LUSHBAUGH
ATTN: E TOMPKINS
ATTN: J TOTTER

OHIO STATE LIBRARY
ATTN: LIBRARIAN

OKLAHOMA DEPT OF LIBS
ATTN: U S GOVT DOCUMENTS

OKLAHOMA, UNIVERSITY OF
ATTN: P ANDERSON

OREGON STATE LIBRARY
ATTN: LIBRARIAN

OREGON, UNIVERSITY OF
ATTN: DOCUMENTS SECTION

OREGON, UNIVERSITY OF
ATTN: B PIROFSKY

PACIFIC NORTHWEST LABORATORY
ATTN: S MARKS

PENNSYLVANIA STATE LIBRARY
ATTN: GOVT PUB SECTION

PENNSYLVANIA UNIV HOSPITAL
ATTN: S BAUM

PENNSYLVANIA, UNIV OF
ATTN: P NOWELL

PEORIA PUBLIC LIBRARY
ATTN: BUS, SCI & TECH DEPT

PHILADELPHIA FREE LIB OF
ATTN: GOVT PUBLICATIONS DEPT

PITTSBURGH, UNIV OF
ATTN: E RADFORD
ATTN: LIBRARY

PITTSBURGH, UNIVERSITY OF
ATTN: N WALD

PUB LIB CINCINNATI & HAMILTON CO
ATTN: LIBRARIAN

RAPID CITY PUBLIC LIBRARY
ATTN: LIBRARIAN

READING PUBLIC LIBRARY
ATTN: LIBRARIAN

RHODE ISLAND LIBRARY, UNIV OF
ATTN: GOVT PUB OFFICE

ROCHESTER UNIV MEDICAL CTR
ATTN: C ODOROFF
ATTN: G CASARETT

ROCHESTER UNIV OF LIB
ATTN: DOCUMENTS SECTION

ROCHESTER, UNIVERSITY OF
ATTN: L HEMPELMANN

SAINT FRANCIS HOSPITAL
ATTN: R BLAISDELL

SAN ANTONIO PUBLIC LIBRARY
ATTN: BUS SCIENCE & TECH DEPT

SCOTTSBLUFF PUBLIC LIBRARY
ATTN: LIBRARIAN

SCRANTON PUBLIC LIBRARY
ATTN: LIBRARIAN

SILAS BRONSON PUBLIC LIBRARY
ATTN: LIBRARIAN

SIMON SCHWOB MEM LIB
ATTN: LIBRARIAN

SIOUX CITY PUBLIC LIBRARY
ATTN: LIBRARIAN

SOUTH CAROLINA, MEDICAL UNIV OF
ATTN: P LIU

DNA-TR-86-120 (DL CONTINUED)

SOUTHEASTERN MASSACHUSETTS UNIV LIB
ATTN: DOCUMENTS SEC

SOUTHERN ALABAMA UNIVERSITY OF
ATTN: LIBRARIAN

SOUTHERN CALIFORNIA UNIV OF
ATTN: J BIRREN

SOUTHERN ILLINOIS UNIVERSITY
ATTN: DOCUMENTS CTR

SOUTHERN MISSISSIPPI UNIV OF
ATTN: LIBRARY

SOUTHERN OREGON COLLEGE
ATTN: LIBRARY

SOUTHERN UTAH STATE COLLEGE LIB
ATTN: DOCUMENTS DEPARTMENT

SOUTHWEST MISSOURI STATE COLLEGE
ATTN: LIBRARY

SPOKANE PUBLIC LIBRARY
ATTN: REFERENCE DEPT

STANFORD UNIV MEDICAL CENTER
ATTN: J BROWN

STANFORD UNIVERSITY
ATTN: L MOSES

STANFORD UNIVERSITY HOSPITAL
ATTN: D DOREMAN

STATE LIBR OF MASS
ATTN: LIBRARIAN

TACOMA PUBLIC LIBRARY
ATTN: LIBRARIAN

TEXAS A & M UNIVERSITY
ATTN: R STONE

TEXAS AT AUSTIN UNIV OF
ATTN: H SUTTON

TEXAS STATE LIBRARY
ATTN: US DOCUMENTS SECTION

TEXAS TECH UNIVERSITY LIBRARY
ATTN: GOVT DOCUMENTS DEPT

TEXAS UNIVERSITY OF
ATTN: C S COOK

TEXAS UNIVERSITY OF
ATTN: R STALLONES

TEXAS UNIVERSITY OF
ATTN: W SUTOW

TEXAS UNIVERSITY OF
ATTN: G TAYLOR

TOLEDO PUBLIC LIBRARY
ATTN: SOCIAL SCIENCE DEPT

TRENTON FREE PUBLIC LIBRARY
ATTN: LIBRARIAN

TULSA UNIVERSITY OF
ATTN: LIBRARIAN

UTAH STATE UNIVERSITY
ATTN: LIBRARIAN

UTAH, UNIVERSITY OF
ATTN: LIBRARY

UTAH, UNIVERSITY OF
ATTN: DOC DIVISION

VANDERBILT UNIVERSITY
ATTN: R QUINN

VERMONT, UNIVERSITY OF
ATTN: DIRECTOR OF LIBRARIES

VIRGINIA STATE LIBRARY
ATTN: SERIALS SECTION

VIRGINIA, UNIVERSITY OF
ATTN: PUBLIC DOCUMENTS

WASHINGTON STATE LIBRARY
ATTN: DOCUMENTS SECTION

WASHINGTON, UNIVERSITY OF
ATTN: D THOMPSON

WASHINGTON, UNIVERSITY OF
ATTN: A MOTULSKY

WEST VIRGINIA, UNIVERSITY OF
ATTN: DIR OF LIB (REGIONAL)

WISCONSIN, UNIVERSITY OF
ATTN: ACQUISITIONS DEPARTMENT

WISCONSIN, UNIVERSITY OF
ATTN: J CROW

WORCESTER PUBLIC LIBRARY
ATTN: LIBRARIAN

YALE UNIVERSITY
ATTN: DIRECTOR OF LIBRARIES

YALE UNIVERSITY SCH OF MEDICINE
ATTN: J MEIGS
ATTN: LIBRARY

END

DATED

FILM

8-88

Dtic